

## **Autologous Osteochondral Bone Grafting (Mosaicplasty)**

### **Introduction**

Surgery: Grafting for cartilage defects of femoral condyles or patello-femoral articulation. Grafts taken from the non-weight bearing surface of the intercondylar notch or the non weight bearing surface of the outer aspect of medial/lateral condyles above the level of the notch. The cartilage defect is drilled and filled with grafts without any cement. Femoral condyle grafting is either arthroscopic or combined arthroscopic and arthrotomy. Patello-femoral grafting is always through arthrotomy.

Indications for surgery: Pain, swelling and mechanical symptoms. These defects can be traumatic, idiopathic, associated with Anterior Cruciate Ligament (ACL) deficient knees or osteochondritis dissecans (OCD).

Expected Length of Stay: Day case

Surgeons: Mr Keating, Mr Lawson, Miss Biant

### **Scope of practice**

These guidelines are designed to guide physiotherapists when treating patients following a Mosaicplasty. These guidelines were produced by a process of systematic review of the current evidence-based literature, medical and peer consultation. They were correct at the time of writing. The guidelines should be used in conjunction with the clinical reasoning skills of the physiotherapist and patients should always be treated on a case-by-case basis.

### **Aim**

The aim of these guidelines is to provide physiotherapy staff with a series of recommendations from the current evidence base to assist them in the management of patients who have undergone a Mosaicplasty.

### **Literature review question**

What is a safe and effective rehabilitation programme for patients undergoing a Mosaicplasty to maximise outcome and return to full activity/sport?

### **Search Process**

Appraisal process: Literature searches using key words were performed on the databases listed below. No articles comparing postoperative management strategies were found but key articles with descriptors of the surgery and rehabilitation were included for information. Of note there are two active trials that are comparing Mosaicplasty to other treatments at the time of update they are not currently published.

Total number of articles selected: 21

Total number of articles discarded: 21

CASPs used: n/a

# LOTHIAN PHYSIOTHERAPY ORTHOPAEDIC GUIDELINES

Data Bases:

Data Base	Dates	Limitations
Medline	Unlimited	None
Cochrane	Unlimited	None
Google	Unlimited	None
Amed	Unlimited	None
Cinahl	Unlimited	None
Embase	Unlimited	None
Sports Discuss	Unlimited	None
Joanna Briggs	Unlimited	None
NHS QIS	Unlimited	None

Key Words:

Mosaicplasty	Post-operative rehabilitation
Osteochondral Autologous Transplantation	Weight-bearing
OATS	Trochlear
Knee surgery	Articular cartilage
Quadriceps exercises	Knee mobilisation
Physiotherapy	

## Results

The literature review revealed no RCT's on physiotherapy management following a Mosaicplasty and therefore, at the time of writing this, no individual article answered the review question in its entirety. Within the available literature there is variation in the post-operative management following a Mosaicplasty. These guidelines therefore reflect current practice and expert opinion and have been developed in consultation with the Orthopaedic Consultants at the Royal Infirmary of Edinburgh.

## Key Points

- ***Please note there are slightly different protocols depending on the anatomical site of grafting***
- Timescales are approximate and rehabilitation should be guided, at each stage, by minimal swelling, resolution of pain and good muscle recruitment

# LOTHIAN PHYSIOTHERAPY ORTHOPAEDIC GUIDELINES

- Patients are usually non-weight bearing (NWB) for the first 6 weeks (Mr Keating's patients can begin partial weight bearing (PWB) at 4 weeks and progress to fully weight bearing (FWB) at 6 weeks)

## **Femoral Condyles**

### **Phase 1**

#### **Immediate Post Operative/Early Rehabilitation: 0-6 weeks**

Goals	Recommendations	
Reduce post-surgical inflammation	Ice 20 minutes, 2 hourly Compress when active Elevate when resting	A1 C
Protect Graft Site	NWB (Mr Keating's patients can PWB at 4 weeks and progress to FWB at 6 weeks, unless otherwise stipulated in post op instructions)	C
Increase knee range of movement (ROM)	Passive/Active-assisted/Active knee flexion & extension	C
Commence quadriceps muscle strengthening	NWB strengthening exercises/open chain quadriceps	C

# LOTHIAN PHYSIOTHERAPY ORTHOPAEDIC GUIDELINES

## Patello-femoral

### Phase 1

#### Immediate Post Operative/Early Rehabilitation: 0-6 weeks

Goals	Recommendations	
Reduce post-surgical inflammation	Ice 20 minutes, 2 hourly Compress when active Elevate when resting	A1 C
Protect Graft Site	NWB (Mr Keating's patients can PWB at 4 weeks and progress to FWB at 6 weeks, unless otherwise stipulated in post op instructions)  No active (open or closed chain) quadriceps exercises  No active range of movement (AROM) exercises over the edge of bed or on a chair (i.e. through range quad) due to increased patellofemoral joint forces	C
Increase knee range of movement (ROM)	Knee flexion/extension on sliding board or prone knee bend only	C
Commence quadriceps muscle strengthening	Static quadriceps contraction only  Hamstring strengthening allowed	C

# LOTHIAN PHYSIOTHERAPY ORTHOPAEDIC GUIDELINES

## **Femoral Condyles and Patello-femoral**

### **Phase 2**

#### **Middle stage: 6-12 weeks**

<b>Goals</b>	<b>Recommendations</b>	
Full range of movement	AROM exercises at the Knee	C
Restore normal gait	Gait re-education as required (FWB allowed)	C
Regain functional lower limb strength	Open and closed chain quadriceps exercises  As required, include strengthening work for other affected muscle groups (i.e. Gluts, Hamstrings, Gastroc/Soleus)	C
Increase neuromuscular control and stability	Single leg balance exercises and dynamic stability work (as weight bearing allows)	C

### **Phase 3**

#### **Late stage: 12 weeks – 6 months**

<b>Goals</b>	<b>Recommendations</b>	
Return to full sport/activity	Sport specific and Plyometric rehabilitation  Ensure full ROM, control and strength before commencing sport specific/plyometric exercises	C

# LOTHIAN PHYSIOTHERAPY ORTHOPAEDIC GUIDELINES

## **Criteria for returning to sport:**

Low impact sports 3-4 months depending on symptoms and progress	C
High impact sports 4-6 months depending on symptoms and progress	C
Non-irritable knee with full ROM	C
Objective measures of ROM, strength and dynamic balance should be recorded	C

## **Potential complications:**

Infection  
Femoral condyle fracture  
Knee stiffness  
Dislodging of grafts (loose bodies)  
Failure to relieve pain  
Patello-femoral pain

If clear signs of complications, refer back to surgeon

## **Expectations of Surgery:**

Pain relief  
Full ROM  
Return to normal activities/sport

## **Contact details for further information/advice:**

Royal Infirmary of Edinburgh, Outpatient Physiotherapy Department: 0131 242 1940

Western General Hospital, Outpatient Physiotherapy Department: 0131 537 1288

St John's Hospital, Outpatient Physiotherapy Department: 01506 522063

# LOTHIAN PHYSIOTHERAPY ORTHOPAEDIC GUIDELINES

## **Bibliography**

Association of Chartered Physiotherapists in Sports and Exercise Medicine (2010). Management of Acute Soft Tissue Injury using Protection, Rest, Ice, Compression and Elevation

Bentley et al (2003) A prospective randomised comparison of ACI versus mosaicplasty for osteochondral defects of the knee, JBJS. 85-B pp223-230  
Buckwalter, J (1998) Articular Cartilage: Injuries and Potential for Healing, J Orthop Sports Phys Ther 28 (4) pp192-202

Dozin et al (2005) Comparative Evaluation of Autologous Chondrocyte Implantation and Mosaicplasty. A Multicentred Randomized Clinical Trial Clin J Sports Med 15(4)

Gudas, R et al (2005) A Prospective Randomised clinical Study of Mosaic osteochondral Autologous Transplantation v Microfracture for the treatment of osteochondral defects in the knee joint in Young Athletes Arthroscopy New York 21(9) pp1066-1075

Hamby, K. et al (2012) Rehabilitation after Articular Cartilage Repair of the Knee in the Football (Soccer) Player. *Cartilage*. 3: 50S-56S

Hangody, L. et al (2010) Clinical Experiences with Autologous Osteochondral Mosaicplasty in an Athletic Population: A 17-year Prospective Multicenter Study. *The American Journal of Sports Medicine*, 38: 1125-1133

Hangody, L and Fules (2003) Autologous Osteochondral Mosaicplasty for the treatment of full thickness defects of weight bearing joints JBJS 85 –A pp25-32

Horas, U et al (2003) Autologous Chondrocyte Implantation and Osteochondral Cylinder Transplantation in Cartilage Repair of the Knee JBJS 85(2) pp185-192

Jones, M and Willams, A (2016). Osteochondritis dissecans of the knee: a practical guide for surgeons. *Bone Joint J* 98-B:723–9

Krych, A. et al (2012) Activity Levels are Higher After Osteochondral Autograft Transfer Mosaicplasty than after Microfracture for Articular Cartilage Defects of the Knee: A Retrospective Comparative Study. *The Journal of Bone & Joint Surgery*, 94: 971-978

Ma H et al (2004) Osteochondral autografts transfer for post traumatic osteochondral defect of the knee – 2-5 years follow up Injury 35 pp1286-1292

Mithoefer, K. et al (2012) Current concepts for Rehabilitation and Return to Sport After Knee Articular Cartilage Repair in the Athlete. *Journal of Orthopaedic & Sports Physical Therapy* 42(3): 254-273

Morelli et al (2002) Management of chondral injuries using osteochondral autogenous transfer (mosaicplasty)

Rimtautas, G. et al (2012) Ten-Year Follow-up of a Prospective, Randomized

## LOTHIAN PHYSIOTHERAPY ORTHOPAEDIC GUIDELINES

Clinical Study of Mosaic Osteochondral Autologous Transplantation Versus Microfracture for the Treatment of Osteochondral Defects in the Knee Joint of Athletes. *The American Journal of Sports Medicine*, 40(11): 2499-2508

Robert, H. (2011) Chondral Repair of the Knee Joint using Mosaicplasty. *Orthopaedics & Traumatology: Surgery & Research* 97: 418-429

Solheim, E. et al (2010) Osteochondral Autografting (Mosaicplasty) in Articular Cartilage Defects in the knee: Results at 5 to 9 years. *The Knee*, 17: 84-87

Solheim, E. et al (2013) Results at 10 to 14 years after Osteochondral Autografting (Mosaicplasty) in Articular Cartilage Defects in the Knee. *The Knee*.

Solheim, E. et al (2018) Randomized Study of Long-term (15-17 Years) Outcome After Microfracture Versus Mosaicplasty in Knee Articular Cartilage Defects. *Am J Sports Med*, 46 (4): 826-831.

Tuluhan, E.Y. et al (2013) Factors affecting the outcome of Osteochondral Autografting (Mosaicplasty) in Articular Cartilage Defects of the knee joint: Retrospective Analysis of 152 cases. *Arch Orthop Trauma Surg*, 133: 531-536

Van Assche D (2007) Cartilage Repair Rehabilitation Issues Position Paper ICRS Rehabilitation Consensus Meeting 29-30 June 07

Wasiak J et al (2007) Autologous cartilage implantation for full thickness articular cartilage defects of the knee (review) Cochrane Collaboration 4 pp1-15



# LOTHIAN PHYSIOTHERAPY ORTHOPAEDIC GUIDELINES

Version: 5

Authors: Jenny Forsyth, Juliet Gold, Elaine Graham

Date of Original: May 2008

Review Author: Molly Wright

Review Date: May 2019

Date for review: 2024

Disclaimer: This information should be used in conjunction with treatment from an appropriate physiotherapist and not in isolation.

## **Appendix**

### **Levels of Evidence**

Evidence from large randomised controlled trials (RCTs) or systematic review (including meta-analyses)† A1

Evidence from at least one high quality cohort A2

Evidence from at least on moderate size RCT or systematic review A3

Evidence from at least one RCT B

Expert opinions C

Laboratory Evidence\* D

† *Arbitrarily, the following cut-off points have been used: large study size \_ 50 patients per intervention group; moderate study size \_ 30 patients per intervention group.*

\* *Arbitrarily, added by Lothian Physiotherapy Musculoskeletal Network Group*

Modified from: MacAuley D and Best TM (2007) Evidence-based Sports Medicine. 2nd Edition. BMJ Books. Blackwell Publishing. Oxford, UK.