

## PHYSIOTHERAPY TREATMENT OPTIONS FOR TENNIS ELBOW

NHS Lothian

### **Introduction:**

Tennis elbow, also known as 'lateral epicondylitis', 'lateral elbow pain' or 'lateral elbow tendinopathy', is a common musculotendinous disorder of the extensor origin at the lateral humeral epicondyle. The condition is considered to be a 'partially reversible but degenerative overuse – underuse tendinopathy' (Orchard and Koutouris, 2011). The question of the pain of tendinopathy, physiological or pathophysiological, remains unanswered; however there is evidence for both – tendon based nociceptive contributions and mechanisms within the periphery and the central nervous system (Rio et al. 2014).

Typical symptoms include lateral elbow pain and tenderness, pain with wrist extension / supination and reduced grip strength. Tennis elbow affects between 1-3% of the general population, with men and women being equally affected (Tosti et al. 2013). It is most prevalent between ages 35-54 years (McMurtie & Watts 2012). There is some evidence of higher incidence of tennis elbow in amateur tennis players and those with manually intensive occupations (Coombes et al. 2008).

### **Aim:**

The aim of this guideline is to provide NHS Lothian physiotherapy staff with options from research findings to assist them in the management of patients with tennis elbow. A previous version of this guideline 'Treatment Options for Tennis Elbow' was written in 2014. This version of the guideline is an updated guideline, written in 2021. An incremental approach to the guideline update was adopted as per the *Cochrane Handbook for Systematic Reviews of Interventions* (Cumpston et al. 2022). This update retains previous text from the 2014 edition and adds new research evidence from studies published between 2014 and 2020.

### **Scope of practice and limitations:**

These treatment options are suggestions to guide physiotherapists when treating patients with tennis elbow. The guidelines are written by a team of physiotherapy clinicians and were produced by a process of review of the research literature and peer consultation. While best efforts were made to include all relevant literature and use comprehensive search methods, the production process has not undergone the same process of academic scrutiny as that of a National Clinical Guideline or a published systematic review. Research included in the guideline is of variable quality and may have a high risk of bias. Any treatment options should be used in conjunction with the clinical reasoning skills of the physiotherapist and patients should always be treated on a case-by-case basis.

### **Literature review question:**

What is the best available evidenced physiotherapy treatment for tennis elbow?

## Literature Search Databases:

### *Original Guideline (2014)*

Database	Dates	Limitations
CINAHL	2006 to 2014	English
Ovid	2006 to 2014	English

### *Updated Version (2021)*

Database	Dates	Limitations
Medline	2014 to 2020	English
SportDiscus	2014 to 2020	English
CINAHL	2014 to 2020	English
Ebsco host	2014 to 2020	English
Cochrane	2014 to 2020	English

## Keywords:

### *Original Guideline (2014)*

Lateral epic*	Physical Therapy
Tennis elbow	Treat
Therapy	Physio
Rehab	Friction
Taping / strapping / bandages / strapping	Anti inflammatory agents
Steroidal Injections	Ultrasonic therapy

### *Updated Version (2021)*

Tennis Elbow or Lateral Epicondylitis or Lateral Tendinopathy or Lateral Elbow Pain	Therapy
Electrotherapy	Management
Massage	NSAIDS or Non-steroidal AIDs
Strength or strengthening	Support
Treatment	Taping or strapping
Intervention	Electrotherapy

## Literature Search Results:

In both the original and updated version of this guideline, research articles were assigned levels of evidence according to 'Sign 50: A guideline developer's handbook'. See reference list for assigned levels.

There was no detail on the criteria for exclusion of studies in the 2014 version of this guideline. In the updated version, only research articles of SIGN level 3 and above were included in the guideline, and those considered relevant to the research question.

Critical Appraisal Skills Programme (CASP) checklists were used to assist with article appraisal.

***Original Guideline (2014)***

Total number of articles in the search	26
Total number of articles discarded	4
Total number of articles reviewed	22

***Updated Version (2021)***

Total number of articles in the search	49
Total number of articles discarded	38
Total number of articles reviewed	11

## **RECOMMENDATIONS:**

### **Summary:**

Many non-operative treatments have been suggested for tennis elbow. Conservative management strategies considered in this guideline include: advice, exercise, manual therapy, electrotherapy, acupuncture, taping and supports, ice, corticosteroid injections and non-steroidal anti-inflammatory drugs (NSAIDs). In both the original guideline (2014) and this updated version (2021), there are wide ranging evidence levels. Details on research and recommendations for each management strategy can be found in the body of this guideline. The mechanism of action for each management strategy is not discussed in this guideline, however, are likely to have an effect both locally at the site of the extensor tendon and on the central nervous system (Rio et al. 2014).

Tennis elbow may also be considered a self-limiting disorder (Shiri and Viikari-Juntura, 2011). Effective conservative treatment generally includes interventions to 'relieve pain, control inflammation, promote healing, improve local and general fitness and control force loads' (Waseem et al. 2012). Patients may improve with non-operative measures (Tosti et al. 2013), while there is also some evidence to suggest that approximately 90% of patients with tennis elbow are symptom free after 12–24 months, some without any intervention (Bisset et al. 2006, Coombes et al. 2013).

### **1. Exercise and Advice**

Consideration should be made regarding the impact of repetitive movements of the hand and wrist, and activities that load the extensor tendons (Shiri & Viikari-Juntura, 2011, Tosti et al. 2013, McMurtie et al. 2011). An ergonomic assessment may be beneficial for patients, and for those whose lifestyle is more sedentary, for example office workers (Shiri & Viikari-Juntura, 2011, Waseem et al. 2012). Several authors recommend examination of daily gripping / pronating / supinating actions and how these may be modified to change the load on the tendons (Frogh et al. 2012, Waseem et al. 2012, Tosti et al. 2013, Shiri & Viikari-Juntura 2011, and De Smedt et al. 2007).

Some research evidence suggests that a rehabilitation program that consists of a mixture of stretching and loading exercise could reduce pain and improve pain-free strength and function in people presenting with tennis elbow (Waseem et al. 2012, Raman et al. 2012, Croisier et al. 2007, Ortega-Castillo & Medina-Porqueres 2016, Stasinopoulos et al. 2010, Keating et al. 2020, Tiwari 2018).

There is mixed evidence as to whether eccentric loading is more beneficial than concentric isometric exercise (Waseem et al. 2012, Raman et al. 2012, Croisier et al. 2007, Ortega-Castillo & Medina-Porqueres 2016).

Exercise programmes in these studies are varied but most commonly consist of three sets of fifteen repetitions, carried out twice daily (Ortega-Castillo & Medina-Porqueres 2016). There is however very little high-quality evidence that conclusively supports this number of repetitions / sets / frequency. One RCT suggested an exercise program carried out three days per week for the first three weeks, followed by six days per week for the remaining three weeks (Lee et al. 2014).

The position of loading assessed in research studies is most frequently with elbow in full extension or flexed, forearm in pronation, arm supported, and wrist extended with hand hanging over the edge (Waseem et al. 2012, Stasinopoulos et al. 2010 and Raman et al. 2012).

Previous guidelines suggested exercising in a pain-free range (Waseem et al. 2012) whereas newer literature from the 2020 search has shown an effective outcome with exercise that allowed for mild discomfort (Keating et al. 2020). Waseem et al. (2012) and Stansinopoulos et al. (2010) suggest three repetitions of a 30-45 second stretch before and after strength training. In these studies, the extensor muscles were stretched with the elbow in full extension, forearm pronated, wrist flexed with ulnar deviation.

## **2. Joint Mobilisations**

Many studies have shown positive effects on pain and grip strength with the application of mobilisations with movement (Amro et al. 2010, Bisset et al. 2006, Vicenzino et al. 2008, Bagade and Verma 2015, Hsu et al. 2016 and Lucado et al. 2018). A small study showed an increase in pain-free grip strength for patient with tennis elbow after T5 joint mobilisation (Zunke et al. 2020), however further research is required to support these isolated findings.

## **3. Soft Tissue Techniques**

The Cochrane review on Deep Transverse Friction Massage for treating Tennis Elbow (Loew et al. 2014) concluded: “We do not have sufficient evidence to determine the effects of deep transverse friction on pain, improvement in grip strength, and functional status for patients with lateral elbow tendinitis as no evidence of clinically important benefits was found”. However, there is mixed evidence to suggest that transverse frictions could be beneficial in the treatment of tennis elbow (Coombes et al. 2008 and Yi, Bratchenko and Tan 2018).

## **4. Taping and Strapping**

The effectiveness of orthotic devices has not been firmly concluded (Coombes et al. 2008). It is suggested that a forearm support reduces the stress around the origin of the wrist extensors at the point of attachment to the epicondyle.

A study by Takasaki et al. (2008) found that positioning the clasp over the proximal 1/5th of the forearm was effective at reducing stress on the extensor carpi radialis brevis. There is low quality evidence to suggest that splinting and taping interventions may have short-term benefits on pain relief and arm function in the management of tennis elbow (Akkurt et al. 2018, Bhambhani et al. 2016, Cho et al. 2018, Guler and Yildirim 2020, Nowotny et al. 2018, Kachanathu et al. 2019, Giray et al. 2019, Shakeri et al. 2018, Vellilappilly et al. 2017). However, this effect on pain, function and strength was relatively short-lived and research suggests that benefits do not extend beyond twelve weeks after treatment (Shahabi et al 2020). This research may not therefore advocate the use of an epiclasp on an ongoing basis. One RCT reported no significant difference between counterforce brace and kinesio-taping at six weeks (Phadk and Desai 2017).

## **5. Acupuncture**

Bisset et al. (2005) conducted a meta-analysis of clinical trials on physical interventions for tennis elbow. In this paper they included four research studies deemed of sufficient methodology quality to be included in the review. The authors conclude that there appears to be some evidence to support the efficacy of acupuncture over placebo as a treatment for tennis elbow pain in the short-term. However, this benefit appears to be short-lived, that is, two to eight weeks.

An audit of acupuncture practice by fourteen 'expert' physiotherapists in managing tennis elbow in 30 patients demonstrated a low level of consistency in practice and application of up to 103 different points (Alvim and Ferreira 2018). The most common of these were LI-11 (297 of 420) and LI-4 (160 of 420). There is no evidence that supports the use of particular acupuncture points in the management of tennis elbow, however a low-quality study by Jeon et al. (2017) suggested that a single treatment of acupuncture (points LI-4, TE-5, LU-5, LI-10, LI-12, SP-6, GB-34) may be effective in providing a small but significant reduction in pain immediately following acupuncture. There was no change in pain-free grip strength and the authors conclude that the use of acupuncture is unlikely to enhance the ability of people with tennis elbow to engage with pain-free rehabilitation.

Studies that examined the use of electroacupuncture, dry point needling and percutaneous tendon needling were excluded from this guideline due to their limited availability to be practised in NHS Lothian.

## **6. Ice**

Several review articles have supported the use of ice packs in tennis elbow as a pain-relieving modality (Waseem et al. 2012, Walz et al. 2010). However there appears to be a lack of research to fully support the use of cryotherapy as a disease-modifying modality.

## **7. Electrotherapy**

There is research comparing Ultrasound and ESWT in treating tennis elbow that shows significant improvement with both techniques, but neither have been shown to be superior to the other (Lizis 2015, Yalvaç et al. 2018, Dedes et al. 2020).

Köksal et al. (2015) suggest Extracorporeal Shockwave Therapy (ESWT) is clinically effective in treating acute and persistent tennis elbow, showing a significant reduction in pain and improvement in function. While Razavipour et al. (2018) show a significant reduction in pain and improvement in Disabilities of the Arm, Shoulder and Hand (DASH) scores at 30 and 60 days after one week of daily application of ESWT. Aydin et al (2018) suggest that ESWT should be used in conjunction with traditional conservative management techniques rather than in isolation.

## **8. Non-Steroidal Anti-Inflammatory Drugs (NSAIDs)**

The use of NSAIDs is not supported in the long term due to potential side-effects and complications (Orchard & Koutouris 2011). However, there is some evidence to support their use in the short term (Shiri & Viikari-Juntura 2011, Waseem et al. 2012).

A Cochrane review on the effects of NSAIDs on tennis elbow (Pattanittum 2013) concluded: "There remains limited evidence from which to draw firm conclusions about the benefits or harms of topical or oral NSAIDs in treating lateral elbow pain. Although data from five placebo-controlled trials suggest that topical NSAIDs may be beneficial in improving pain (for up to four weeks), non-normal distribution of data and other methodological issues precluded firm conclusions. Some people may expect a mild transient skin rash. Evidence about the benefits of oral NSAIDs has been conflicting, although oral NSAID use may result in gastrointestinal adverse events in some people. No direct comparisons between oral and topical NSAIDs were available. Some trials demonstrated greater benefit from glucocorticoid injection than from NSAIDs in the short term, but this was not apparent in all studies and was not apparent by 6 months in the only study that included longer-term outcomes."

## **9. Cortico-Steroid Injections**

Moderate evidence exists which supports the use of Cortico-Steroid Injections (CSI) for short-term pain relief and improvement in self-reported function (Snyder and Todd 2012, Coombes et al. 2013, Beyazal & Devrimsel 2015, Barnett et al. 2019, Xiong et al. 2019).

Lin-Fen Hsieh et al. (2018) suggests there is no short-term significant difference between a CSI and lidocaine injection for Tennis Elbow.

Long-term studies by Bisset et al. (2006) and Coombes et al. (2013) report worse clinical outcomes and a higher incidence of recurrence of symptoms when CSI is the initial line of treatment. Regenerative injections and ESWT have been shown to have significant results compared with CSI up to two years post-treatment (Barnett et al. 2019 and Xiong et al.2019).

Coombes et al. (2015) demonstrates that CSI is associated with great variability, and a low probability of being cost-effective. They concluded that physiotherapy should be used as first line intervention before considering CSI.

## **10. Surgery**

Less than 5% of patients fail to respond to conservative treatment (Tosti et al. 2013). Waseem et al (2012) suggest that surgery may be considered if conservative therapy has been unsuccessful over prolonged periods in those people where there is evidence of extra-articular calcification or there are multiple areas of tendonitis.

## References

Level of Evidence	Reference
1+	Ajimsha, M.S., Chithra, S. and Thulasyammal, RP. 2012 Effectiveness of Myofascial Release in the Management of Lateral Epicondylitis in Computer Professionals. Archives of Physical Medical Rehabilitation. Vol 93. pp 604-609
1+	Akin, C., Okin, O., and Koseoglu, BF. 2010. Short-term Effectiveness of Ultrasound Treatment in Patients with Lateral Epicondylitis: Randomized, Single-blind, Placebo-controlled, Prospective Study. Turkish Journal of Rheumatology. 25 (2) pp. 50-55
2+	Akkurt, H.E. <i>et al.</i> 2018. Comparison of an epicondylitis bandage with a wrist orthosis in patients with lateral epicondylitis. Prosthetics and orthotics international. 42(6), pp. 599–605. doi:10.1177/0309364618774193.
2+	Amro, A., Diner, I., Badair, WO., Hamada, IM., Shalabi, AI., and Ilyyan, DI. 2010. The Effects of Mulligan Mobilisation with Movement and Taping Techniques on Pain, Grip Strength, and Function in Patients with Lateral Epicondylitis. Hong Kong Physiotherapy Journal. 28 pp. 19-23
2+	Alvim, D. and Ferreira, A. 2018. Pragmatic Combination of Acupuncture Points for Lateral Epicondylalgia are Unreliable in the Physiotherapy Setting. Journal of Acupuncture and Meridian Studies. 11(6) pp. 367-374.
1-	Aydın, C.G., Aykut, S., Arslanoğlu, F., Kılınc, C.Y. and Kocaer, N., 2017. Long-term efficiency of extracorporeal shockwave therapy on lateral epicondylitis. 83 pp. 438-444
1-	Bagade, V.K. and Verma, C., 2015. Effect of Mulligan Mobilization with Movement (MWM) in the Treatment of Chronic Lateral Epicondylitis: 24 Weeks Follow-up Study. Website: www. ijpot. com, 9(4), p.199.
1+	Barnett, J., Bernacki, M.N., Kainer, J.L., Smith, H.N., Zaharoff, A.M. and Subramanian, S.K., 2019. The effects of regenerative injection therapy compared to corticosteroids for the treatment of lateral Epicondylitis: a systematic review and meta-analysis. Archives of physiotherapy, 9(1), pp.1-12.
1+	Beyazal, M.S. and Devrimsel, G., 2015. Comparison of the effectiveness of local corticosteroid injection and extracorporeal shock wave therapy in patients with lateral epicondylitis. Journal of physical therapy science, 27(12), pp.3755-3758.
1-	Bhambhani, S., Mitra, M. and Kaur, A. 2016. Effectiveness of Kinesiotaping along with Conventional Physiotherapy for Patients with Tennis Elbow. Indian Journal of Physiotherapy and Occupational Therapy. 10(3) pp. 18-23.
1++	Bisset, L., Beller, E. Jull, G. Brooks, P, Darnell R and Vicenzio, B. 2006. Mobilisation with Movement and Exercise, Corticosteroid Injection, or Wait and See for Tennis Elbow: Randomised Trial. BMJ. 333 pp. 939-941
1++	Bisset, L., Paungmali, A., Vicenzino, B., et al 2005. A systematic review and meta-analysis of clinical trials on physical interventions for lateral epicondylalgia. British Journal of Sports Medicine. 39 pp. 411-422.
2++	Cho, Y.T. et al 2018. Kinesio taping reduces elbow pain during resisted wrist extension in patients with chronic lateral epicondylitis: a randomized, double-blinded, cross-over study. BMC musculoskeletal disorders. 19(1), pp. 193–193. doi:10.1186/s12891-018-2118-3.
4	Coombes, B.K., Bisset, L. and Vicenzino, B. 2008. A new integrative model of lateral epicondylalgia. British Journal of Sports Medicine. 43 (4) pp: 252-8
1-	Coombes, B.K., Bisset L., Brooks P., Kahn, A. and Vicenzio B. 2013. Effect of Corticosteroid Injection, Physiotherapy or both on clinical Outcomes in Patients With Unilateral Lateral Epicondylalgia. A Randomized Controlled Trial. Journal of the American Medical Association. 309 (5) pp. 461 – 469
1-	Coombes, B.K., Connelly, L., Bisset, L. and Vicenzino, B., 2016. Economic evaluation favours physiotherapy but not corticosteroid injection as a first-line intervention for chronic lateral epicondylalgia: evidence from a randomised clinical trial. British journal of sports medicine, 50(22), pp.1400-1405.
2++	Croisier, J., Dessalle-Foidart, M., Tinant, F., Crielaard, J.M., Forthomme, B. 2007. An Isokinetic Eccentric



	Programme for the Management of Chronic Lateral Epicondylar Tendinopathy. <i>British Journal of Sports Medicine</i> . 41 (4) pp. 269-275
N/A	Cumpston M, Chandler J. Chapter IV: Updating a review. In: Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, Welch VA (editors). <i>Cochrane Handbook for Systematic Reviews of Interventions</i> version 6.3 (updated February 2022). Cochrane, 2022. Available from <a href="http://www.training.cochrane.org/handbook">www.training.cochrane.org/handbook</a> .
4	De Smedt, T., De Jong, A., Leemput, W.V., Lieven, D., and Van Glabbeek, F. 2007. Lateral Epicondylitis in Tennis: Update on Aetiology, Biomechanics and Treatment. <i>British Journal of Sports Medicine</i> . 41 (11) pp. 816-819
1-	Dedes, V., Tzirogiannis, K., Polikandrioti, M., Dede, A.M., Mitseas, A. and Panoutsopoulos, G.I., 2020. Comparison of radial extracorporeal shockwave therapy with ultrasound therapy in patients with lateral epicondylitis. <i>Journal of Medical Ultrasonics</i> , 47(2), pp.319-325.
1-	Forogh, B. Kahalahi, M., Javamashir, MA., Ghoseiri, K., Kamali, M. and Raissi, G. 2012. The Effects of a New Designed Forearm Orthosis in Treatment of Lateral Epicondylitis. <i>Disability &amp; Rehabilitation: Assistive Technology</i> . 7 (4) pp. 336 -339
2+	Giray, E, Karali-Bingul, D, Akyuz, G. 2019. The Effectiveness of Kinesiotaping, Sham Taping or Exercises Only in Lateral Epicondylitis Treatment: A Randomized Controlled Study. <i>Physical Medicine and Rehabilitation</i> 11 pp. 681-693.
3	Gonzalez-Iglesias, J., Cleland, J. del Rosario, M. and Fernandez de las Penas, F. 2011. Case Series. Multimodal Management of Lateral Epicondylalgia in Rock Climbers: A Prospective Case Series. <i>Journal of Manipulative and Physiological Therapeutics</i> . 34 (9) pp.635-641
2+	Guler, T. and Yildirim, P. 2020. Comparison of the Efficacy of Kinesiotaping and Extracorporeal Shock Wave Therapy in Patients with Newly Diagnosed Lateral Epicondylitis: A Prospective Randomized Trial. <i>Nigerian Journal of Clinical Practice</i> . 23 pp. 704-710.
1++	Hsieh, L.F., Kuo, Y.C., Lee, C.C., Liu, Y.F., Liu, Y.C. and Huang, V., 2018. Comparison between corticosteroid and lidocaine injection in the treatment of tennis elbow: a randomized, double-blinded, controlled trial. <i>American journal of physical medicine &amp; rehabilitation</i> , 97(2), pp.83-89.
1+	Hsu, C.Y., Lee, K.H., Huang, H.C., Chang, Z.Y., Chen, H.Y. and Yang, T.H., 2016. Manipulation therapy relieved pain more rapidly than acupuncture among lateral epicondylalgia (tennis elbow) patients: a randomized controlled trial with 8-week follow-up. <i>Evidence-Based Complementary and Alternative Medicine</i> , doi: 10.1155/2016/3079247
2-	Jeon, J., Bussin, E. and Scott, A 2017. Temporal divergence of changes in pain and pain-free grip strength after manual acupuncture or electroacupuncture: an experimental study in people with lateral epicondylalgia. <i>Chinese Medicine</i> 12(22) pp. 2-11.
2+	Kachanathu, S., Alenazi, A., Hafex, A., Algarni, A. and Alsubiheen, A. 2019. Comparison of the effects of short-duration wrist joint splinting combined with physical therapy and physical therapy alone on the management of patients with lateral epicondylitis. <i>European Journal of Physical and Rehabilitation Medicine</i> . 55(4) pp. 488-93.
1-	Keating, C., Bodnar, R., Joseph, J., Knapp, S., Lepage, M., Solger, G. 2020. Effectiveness of Painful Loading in Lateral Elbow Tendinopathy on Pain Outcomes: A systematic literature review. <i>Orthopaedic Practice</i> . 32 (4) pp. 208-214.
2+	Koksal, I., Guler, O., Mahirogullari, M., Mutlu, S., Çakmak, S. and Aksahin, E., 2015. Comparison of extracorporeal shock wave therapy (ESWT) in acute and chronic lateral epicondylitis. <i>Acta orthopaedica et traumatologica turcica</i> , 49(5), p.465.
1-	Lee, S., Ko, Y., Lee, W. 2014. Changes in Pain, Dysfunction, and Grip Strength of Patients with Acute Lateral Epicondylitis Caused by Frequency of Physical Therapy: A Randomized Controlled Trial. <i>Journal Physical Therapy Science</i> . 26 pp. 1037-1040.
1-	Lizis, P., 2015. Analgesic effect of extracorporeal shock wave therapy versus ultrasound therapy in chronic tennis elbow. <i>Journal of physical therapy science</i> , 27(8), pp.2563-2567.
1++	Loew, L.M., Brosseau, L., Tugwell, P., Wells, G.A., Welch, V., Shea, B., Poitras, S., De Angelis, G. and Rahman, P., 2014. Deep transverse friction massage for treating lateral elbow or lateral knee tendinitis. <i>Cochrane Database of Systematic Reviews</i> , (11).

1++	Lucado, A.M., Dale, R.B., Vincent, J. and Day, J.M., 2019. Do joint mobilizations assist in the recovery of lateral elbow tendinopathy? A systematic review and meta-analysis. <i>Journal of Hand Therapy</i> , 32(2), pp.262-276.
3	Marcolino, A. Siqueira das Neves, L. Oliverira, B., Alexandre, A., Corsatto, G., Barbosa, R., Fonseca, M. 2016. Multimodal approach to rehabilitation of the patients with lateral epicondylitis: a case series. <i>Springerplus</i> . 5 pp. 1718.
1++	McMurtie, A. and Watts, A.C. 2012. Tennis Elbow and Golfer's Elbow. <i>Orthopaedics and Trauma</i> . 26 (5) pp. 337-344
2++	Nowotny, El-Zayat, B., Goronzy, J., Biewener, A., Bausenhardt, F., Greiner, S., & Kasten, P. 2018. Prospective randomized controlled trial in the treatment of lateral epicondylitis with a new dynamic wrist orthosis. <i>European Journal of Medical Research</i> . 23(1) pp. 43-43. <a href="https://doi.org/10.1186/s40001-018-0342-9">https://doi.org/10.1186/s40001-018-0342-9</a>
1+	Ortega-Castillo, M., Medina-Porqueres, I. 2016. Effectiveness of the eccentric exercise therapy in physically active adults with symptomatic shoulder impingement or lateral epicondylar tendinopathy: A systematic review. <i>Journal of Science and Medicine</i> . 19 pp. 438-453.
2++	Orchard, J. and Kountouris, A. 2011. The Management of Tennis Elbow. <i>BMJ</i> . 342 (7812) pp. 1377 - 1382
1++	Pattanittum P, Turner T, Green S, Buchbinder R. Non-steroidal anti-inflammatory drugs (NSAIDs) for treating lateral elbow pain in adults. <i>Cochrane Database of Systematic Reviews</i> 2013, Issue 5. Art. No.: CD003686. DOI: 10.1002/14651858.CD003686.pub2
2-	Phadke, S. and Desai, S. 2017. Effectiveness of Kinesiotape Versus Counterforce Brace as an Adjunct to Occupational Therapy in Lateral Epicondylitis. <i>Indian Journal of Physiotherapy and Occupational Therapy</i> . 11(2). pp. 42-47.
2+	Raman, J., MacDermid, J. And Grewal, R. 2012. Effectiveness of Different Methods of Resistance Exercise in Lateral Epicondylitis-A Systematic Review. <i>Journal of Hand Therapy</i> . 25 pp. 5-26
2-	Razavipour, M., Azar, M.S., Kariminasab, M.H., Gaffari, S. and Fazli, M., 2018. The short term effects of shock-wave therapy for tennis elbow: a clinical trial study. <i>Acta Informatica Medica</i> , 26(1), p.54.
1+	Ruane, H., Hay, L., and Callaghan, M. 2010. Best Evidence Topic. BET 2. Is electrotherapy useful for tennis elbow? <i>Emergency Medicine Journal</i> . 27 (2) pp142-4
1+	Rio, E., Moseley, L., Purdam, C., Smirix, T., Kidgell, D., Pearce, A., Jaberzadeh, S. And Cook, J. 2014. The Pain of Tendinopathy: Physiological or Pathophysiological? <i>Sports Medicine</i> . 44: pp. 9-23.
1+	Shahabi, Bagheri Lankarani, K., Heydari, S. T., Jalali, M., Ghahramani, S., Kamyab, M., Tabrizi, R., & Hosseinabadi, M. 2020. The effects of counterforce brace on pain in subjects with lateral elbow tendinopathy: A systematic review and meta-analysis of randomized controlled trials. <i>Prosthetics and Orthotics International</i> , 44(5) pp. 341-354. <a href="https://doi.org/10.1177/0309364620930618">https://doi.org/10.1177/0309364620930618</a>
2+	Shakeri, H., Soleimanifar, P., Arab, P. and Behbahani, P. 2018. The effects of KinesioTape on the treatment of lateral epicondylitis. <i>Journal of Hand Therapy</i> . 31 pp. 35-41.
1+	Shiri, R., Viikari – Junctura, E. 2011. Lateral and Medial Epicondylitis: Role of Occupational Factors. <i>Balliere's Best Practice &amp; Research in Clinical Rheumatology</i> 25 (1) pp. 43-57
1+	Snyder, K., and Evans, T. 2012. Effectiveness of Corticosteroids in the Treatment of Lateral Epicondylitis. <i>Journal of sports Rehabilitation</i> . 21 (1) pp.83-8.
2++	Stasinopoulos, D., Johnson, M. 2007. It may be Time to Modify the Cyriax Treatment of Lateral Epicondylitis. <i>Journal of Bodywork &amp; Movement Therapies</i> 11 (1) pp. 64-67
2+	Stasinopoulos, D., Stasinopoulos, I., Pantelis, M., and Stasinopolu, K. 2010. Comparison of effects of a home exercise programme and a supervised exercise programme for the management of lateral elbow tendinopathy. <i>British Journal of Sports Medicine</i> . 44 (8) pp. 579-83
1+	Staples, M. Forbes, A. Ptasznik, R., Gordan, J and Buchbinder, R. 2008. A Randomised Controlled Trail of Extracorporeal Shock Wave Therapy for Lateral Epicondylitis (Tennis Elbow). <i>Journal of Rheumatology</i> . 35 (10) pp. 2038-2045
2+	Takasaki, H. Aoki, M., Oshiro, S., Izumi, T. Hidika, E., Fugii, M. and Tatsumi, H. 2008. Strain Reduction of the Extensor Carpi Radialis Brevis Tendon Proximal Origin Following the Application of a Forearm Support Band. <i>Journal of Orthopaedic &amp; Sports Physical Therapy</i> . 38 (5) pp. 257-261

2-	Tiwari, M. 2018. Effectiveness of Flex Bar Eccentric Exercises Versus Progressive Resistance Exercises of Wrist on Pain, Pain free grip strength and Functional Activities in Sub Acute Tennis Elbow. Indian Journal of Physiotherapy and Occupational Therapy. 12(3) 28-32.
2++	Tosti, R. Jennings, J. and Sowards, M. 2013. Lateral Epicondylitis of the Elbow. American Journal of Medicine. 126 (4) pp. 357-e1-357
1-	Vellilappilly, D., Rai, H., Varghese, J. and Renjith, V. 2017. Counterforce orthosis in the management of lateral epicondylitis. Journal Ayub Med Coll 29(2) pp. 328-334.
1+	Vicenzino, B., Smith D., Clelland, J. and Bisset L. 2009. Development of a clinical prediction rule to identify initial 1+ responders to mobilisation with movement and exercise for lateral epicondylalgia. Manual Therapy 14(5) pp 550-4
1-	Viswas, R., Ramachandran, R., and Korde Anatkumar, P. 2012. Comparison of effectiveness of supervised exercise program and cyriax physiotherapy in patients with tennis elbow (lateral epicondylitis) a randomised clinical trial. The Scientific World Journal. Article ID 939645
4	Walz, D., Newman, J.S., Konin, G.P. and Ross G. 2010. Epicondylitis: Pathogenesis, Imaging and Treatment. RadioGraphics. 30 pp167 -184
2++	Waseem, M., Nuhmani, S. and Sachin, Y. 2012. Lateral Epicondylitis: A Review of the Literature. Journal of Musculoskeletal Rehabilitation. 25 (2) pp. 131-142
1+	Xiong, Y., Xue, H., Zhou, W., Sun, Y., Liu, Y., Wu, Q., Liu, J., Hu, L., Panayi, A.C., Chen, L. and Yan, C., 2019. Shock-wave therapy versus corticosteroid injection on lateral epicondylitis: a meta-analysis of randomized controlled trials. The Physician and sportsmedicine, 47(3), pp.284-289.
1+	Yalvaç, B., Mesci, N., Külcü, D.G. and Yurdakul, O.V., 2018. Comparison of ultrasound and extracorporeal shock wave therapy in lateral epicondylitis. Acta orthopaedica et traumatologica turcica, 52(5), pp.357-362.
2++	Yi, R., Bratchenko, W.W. and Tan, V., 2018. Deep friction massage versus steroid injection in the treatment of lateral epicondylitis. Hand, 13(1), pp.56-59.
1+	Zunke, P., Auffarth, A., Hitzl, W. and Moursy, M., 2020. The effect of manual therapy to the thoracic spine on pain-free grip and sympathetic activity in patients with lateral epicondylalgia humeri. A randomized, sample sized planned, placebo-controlled, patient-blinded monocentric trial. BMC musculoskeletal disorders, 21(1), pp.1-11.

**Key: Levels of Evidence from: 'SIGN 50: A guideline developer's handbook'.**

1++ High quality meta-analyses, systematic reviews of RCTs, or RCTs with a very low risk of bias
1+ Well conducted meta-analyses, systematic reviews, or RCTs with a low risk of bias
1-Meta-analyses, systematic reviews, or RCTs with a high risk of bias
2++ High quality systematic reviews of case control or cohort studies. High quality case control or cohort studies with a very low risk of confounding or bias and a high probability that the relationship is causal
2+ Well conducted case control or cohort studies with a low risk of confounding or bias and a moderate probability that the relationship is causal
2-Case control or cohort studies with a high risk of confounding or bias and a significant risk that the relationship is not causal
3 Non-analytic studies, eg case reports, case series
4 Expert opinion